CITY OF DERBY ACADEMY

Farmhouse Road, Sinfin, Derby DE24 3AR

Tel: 01332 270 450 Email: admin@cityofderbyacademy.org

Web: www.cityofderbyacademy.org Headteacher: Mr A Harding BA (Hons) PGCE



December 2022

Dear Parent/Carer

Re: Scarlet fever / Invasive Group A Strep (iGAS)

The UK Health Security Agency (UKHSA) has reported that there is a national increase of scarlet fever above normal seasonal levels. This letter provides information that is relevant for your child and younger children in your family.

Currently, there are no reported cases of either scarlet fever or strep at CODA.

Signs and symptoms of scarlet fever

Scarlet fever is a common childhood infection. It is not usually serious but should be treated with antibiotics to reduce the risk of complications (such as pneumonia) and spread to others. The early symptoms of scarlet fever include sore throat, headache, fever, nausea and vomiting. After 12 to 48 hours, the characteristic red, pinhead rash develops, typically first appearing on the chest and stomach, then rapidly spreading to other parts of the body, and giving the skin a sandpaper-like texture. The scarlet rash may be harder to spot on darker skin, although the 'sandpaper' feel should be present. Patients typically have flushed cheeks and will be pale around the mouth. This may be accompanied by a bright red 'strawberry' tongue.

If you think you, or your child, might have scarlet fever:

- Contact your GP or NHS 111 as soon as possible
- Make sure that you or your child take(s) the full course of any antibiotics prescribed. Although you or your child
 will feel better soon after starting the course of antibiotics, you must complete the course to ensure that you do
 not carry the bacteria in your throat after you have recovered
- Stay at home, away from nursery, school or work for at least 24 hours after starting the antibiotic treatment, to avoid spreading the infection

You can help stop the spread of infection through frequent hand washing and by not sharing eating utensils, clothes, bedding and towels. All contaminated tissues should be disposed of immediately.





Invasive Group A Strep (iGAS)

The same bacteria which cause scarlet fever can also cause a range of other types of infection such as skin infections

(impetigo) and sore throat. In very rare cases, the bacteria can get into the bloodstream and cause an illness called

invasive group A strep (iGAS). Whilst still very uncommon, there has been an increase in iGAS cases this year, particularly

in children under 10 years old.

As a parent, you should trust your own judgement.

Contact NHS 111 or your GP if:

• Your child is getting worse

Your child is feeding or eating much less than normal

• Your child has had a dry nappy for 12 hours or more or shows other signs of dehydration

Your baby is under 3 months and has a temperature of 38C, or is older than 3 months and has a temperature of

39C or higher

Your baby feels hotter than usual when you touch their back or chest, or feels sweaty

Your child is very tired or irritable

Call 999 or go to A&E if:

• Your child is having difficulty breathing – you may notice grunting noises or their tummy sucking under their ribs

• There are pauses when your child breathes

Your child's skin, tongue or lips are blue

• Your child is floppy and will not wake up or stay awake

Stop the spread

During periods of high incidence of scarlet fever, there may also be an increase in outbreaks in schools, nurseries and

other childcare settings. Children and adults with suspected scarlet fever should stay off nursery / school / work until 24

hours after the start of appropriate antibiotic treatment. Good hygiene practice such as hand washing remains the most

important step in preventing and controlling spread of infection.

Yours sincerely

Adrian Harding Headteacher

City of Derby Academy